


U.S. Geological Survey (USGS) Maryland-Delaware-District of Columbia Monthly Water Conditions Summary

USGS June 2018 Water Conditions Summary

Hydrologic data from 28 wells and 33 streamgages were used to monitor the monthly groundwater and streamflow response to weather conditions in Maryland, Delaware, and the District of Columbia. In June 2018, groundwater levels and monthly mean streamflows were at record June highs at one well and two streamgages. Groundwater levels were normal to above normal in 93 percent of the wells and 100 percent of the streams.

The number of groundwater and surface-water sites, and the percentage of the total number of groundwater or surface-water sites, that fall within each percentile range are shown in the graphic for June 2018.



A **percentile** is a value on a scale from 0 to 100 that indicates the percent of a distribution that is equal to or below it. A percentile between 25 and 75 is considered normal. For example, a groundwater level in the 90th percentile is equal to or greater than 90 percent of the values recorded for that month.

Why is it important for the USGS to collect and analyze water-resources data?

USGS water data are valuable to the public, researchers, water managers, planners, and agricultural users, especially during extreme conditions like floods and droughts. The USGS is known for its consistent measurement techniques and the most extensive set of historical groundwater and streamflow data available to the public. Since these long-term data were collected during wet and dry periods, they can be used to assess how water resources respond to changes in weather, and to evaluate how current data compare to the historical data. The uniformity of the dataset enables multi-state comparisons and other comparative statistical analyses that can better inform policy makers of possible water-resources conditions they might encounter in the future.

The sites used in this water summary were carefully selected to include long-term datasets and show the response of streamflow and groundwater levels to weather conditions, rather than the effects of human influences. Of the USGS sites presented in this summary, 13 wells and 29 streamgages have more than 50 years of data. The current streamflow and groundwater data are ranked in comparison to the historical record and summarized. In addition to groundwater and streamflow data, this summary includes precipitation data to give a more complete picture of the region's water resources. Hydrologic and weather data have not been reviewed and are therefore provisional and subject to revision.


U.S. Geological Survey (USGS) Maryland-Delaware-District of Columbia Monthly Water Conditions Summary

Precipitation

Monthly data from five Mid-Atlantic National Weather Service (NWS) stations are used to present monthly precipitation to compare to the response of groundwater and streamflow. The NWS uses averages of data over the 30-year climate normal period from 1981 through 2010.


June 2018 Precipitation

The long-term average June precipitation for the five Mid-Atlantic NWS weather stations is shown in the graph and map.



National Weather Service Stations

- Baltimore =**
Baltimore/Washington International Thurgood Marshall Airport (BWI)
- Georgetown =**
Georgetown, Sussex County Airport
- Hagerstown =**
Hagerstown Regional Airport
- Arlington =**
Ronald Reagan Washington National Airport
- Wilmington =**
New Castle Airport



Source: National Weather Service
DC and MD: <http://w2.weather.gov/climate/index.php?wfo=lwx>
DE: <http://www.weather.gov/climate/index.php?wfo=phi>
MARFC: http://www.weather.gov/marfc/Precipitation_Departures

U.S. Geological Survey (USGS) Maryland-Delaware-District of Columbia Monthly Water Conditions Summary

Groundwater

The USGS monitors groundwater levels in surficial or unconfined aquifers, providing observations that can be compared to both short-term and long-term changes in weather conditions. The groundwater wells used for the monthly water summary were selected based on the following criteria:


- Located in a surficial or unconfined (water-table) aquifer
- Open to a single, known hydrogeologic unit/aquifer
- Groundwater hydrograph generally reflects response to weather
- No indicated nearby pumpage, and likely to remain uninfluenced by pumpage or changes related to human activities
- Minimum period of record is 10 years of continuous/monthly records
- Minimally affected by irrigation, canals, drains, pipelines, and other potential sources of artificial recharge
- Well has a casing – dug wells are generally not used
- Water levels show no apparent hydrologic connection to nearby streams
- Well rarely goes dry
- Long-term accessibility likely, such as on public land

In the Maryland, Delaware, and District of Columbia region, it is useful to compare current data to historical data, such as data from the droughts of 1999-2002 and the 1960s. There are 11 wells that have over 60 years of groundwater data, and 23 wells that have more than 30 years of groundwater data as of 2018.

June 2018 Groundwater Levels

Thirty-nine percent, or 11 USGS observation wells, had groundwater levels within the normal range (25th-75th percentiles) in June. Groundwater levels were above normal at 15 wells, including 1 record high June groundwater level, 5 wells with groundwater levels greater than the 90th percentile, and 9 wells in the 76th-90th percentile range.

There were two wells with below normal groundwater levels. Between May and June, groundwater levels decreased at 14 wells (50 percent) and increased at 14 wells (50 percent).




To access the clickable groundwater map, go to:
http://md.water.usgs.gov/groundwater/web_wells/current/water_table/counties/


U.S. Geological Survey (USGS) Maryland-Delaware-District of Columbia Monthly Water Conditions Summary

In the 5-year hydrographs for the selected wells, groundwater levels are shown as a dark blue line. Each monthly measurement is colored according to the percentile rank compared to the historical values at the site for the month. The normal range is displayed as a white band and is based on the period of record. The maximum water level is at the top of the upper blue section, and the minimum water level is at the bottom of the lower blue area in the graph.

The groundwater level at observation well BA Dc 444, in Baltimore County, Maryland was 39.52 feet below land surface. Normal June groundwater levels at this well range from 36.38 to 39.35 feet below land surface. Groundwater levels have been below normal for the past 15 months. Monthly record-keeping at this well began in September 1988.



The groundwater level at USGS observation well WA Be 2, in Washington County, Maryland, was 15.71 feet below land surface, which is a record June high. The previous record of 19.30 feet below land surface was set in 2003. The normal range of groundwater levels for June at this well is between 27.81 and 33.13 feet below land surface. Record-keeping at this well began in December 1949.



Five-year groundwater hydrographs can be viewed at:
http://md.water.usgs.gov/groundwater/web_wells/current/water_table/counties

U.S. Geological Survey (USGS) Maryland-Delaware-District of Columbia Monthly Water Conditions Summary

Streamflow

Streamflow data are most commonly used for assessing water supply and to determine the risk of droughts and floods. Streamflow data are also used to calculate loads of chemical constituents, and to assess how biological communities are affected by hydrologic conditions.

The USGS streamgages chosen for the monthly water summary were selected based on the following criteria:


- Minimum period of record is 10 years of continuous data
- Watershed areas greater than 5 square miles
- Streamflow is not regulated, such as by a dam or diversion, and has relatively natural flow
- Streamflow data reflect a response to weather conditions
- Most of the surrounding area and watershed are not urban

Of the 33 streamgages used in this summary, 22 have more than 60 years of data, allowing for comparison to data from the historical droughts of 1999--2002 and the 1960s. All 33 streamgages have at least 30 years of monthly mean streamflow data.

June 2018 Streamflow

Monthly mean streamflows were in the normal range (25th-75th percentiles) at 15 percent, or 5 of 33 selected USGS streamgages. Streamflow was above normal at the remaining 28 streamgages including 2 record high June monthly mean streamflows, 18 streamgages with streamflow above the 90th percentile, and 8 streamgages with streamflow between the 76th-90th percentiles.

Streamflow decreased at 25 streamgages (76 percent) and increased at 8 streamgages (24 percent) between May and June.




To access the clickable streamflow map, go to:
<http://md.water.usgs.gov/surfacewater/streamflow/>


U.S. Geological Survey (USGS) Maryland-Delaware-District of Columbia Monthly Water Conditions Summary

In the hydrograph for the selected streamgages, the dark line in the 5-year hydrograph represents the monthly mean streamflow for this period, and the white band shows the normal range (25th-75th percentiles) based on the period of record. The maximum monthly mean streamflow is at the top of the blue shaded section, and the lowest monthly mean streamflow is at the bottom of the tan area. Each monthly mean streamflow is colored according to the percentile rank compared to the historical data for the month.

Monthly mean streamflow at Morgan Creek near Kennedyville in Kent County, Maryland, was 8.82 cubic feet per second (ft³/s). The normal streamflow range for June is between 5.67 ft³/s and 12.40 ft³/s. Record-keeping at this streamgage began in June 1951.



At Potomac River at Hancock, in Washington County, Maryland, the monthly mean streamflow was 14,900 ft³/s, which is a record high. The previous record high was 13,400 ft³/s in 1972. Record-keeping at this streamgage began in October 1932.



Five-year hydrographs can be viewed at:
<http://md.water.usgs.gov/surfacewater/streamflow/>